

[Go to Product page](#)

Datasheet for ABIN1703870
anti-CLCNKB antibody (AA 51-150) (Cy5)

Overview

Quantity:	100 µL
Target:	CLCNKB
Binding Specificity:	AA 51-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CLCNKB antibody is conjugated to Cy5
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CLCNKB
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	CLCNKB
Alternative Name:	CLCNKB (CLCNKB Products)
Background:	Synonyms: Bartter syndrome type 3, Chloride channel Kb, Chloride channel kidney B, Chloride

Target Details

channel protein CLC-Kb, Chloride channel voltage sensitive Kb, CLC K2, CLC-K2, CLCK2, CLCKB, CLCKB_HUMAN, CLCNKB, hCLC Kb, hCLCKb, MGC24087, OTTHUMP00000011120, OTTHUMP00000011121, RP11 5P18.8.

Background: The family of voltage-dependent chloride channels (CLCs) regulate cellular trafficking of chloride ions, a critical component of all living cells. CLCs regulate excitability in muscle and nerve cells, aid in organic solute transport, and maintain cellular volume. CLC-KA is a kidney-specific chloride channel that mediates transepithelial chloride transport in the thin ascending limb of the Henle loop in the inner medulla. CLC-KA plays a crucial role in urine concentration. The gene encoding human CLC-KA maps to chromosome 1p36. Mutations in this gene may be associated with nephrogenic diabetes insipidus in those cases where mutations in the vasopressin V2 receptor and the AQP2 water channel are lacking. CLC-KB mediates basolateral chloride ion efflux in the thick ascending limb and in more distal nephron segments. The gene encoding human CLC-KB maps to chromosome 1p36. Mutations in this gene cause type III Bartter's syndrome which is characterized by renal salt-wasting and low blood pressure.

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months