antibodies - online.com







anti-DPP10 antibody (AA 631-730) (Cy5)



()	ve	K\ /		A .
	\cup	1 V/	-	V۷

Quantity:	100 μL
Target:	DPP10
Binding Specificity:	AA 631-730
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DPP10 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DPP10	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Horse	
Purification:	Purified by Protein A.	

Target Details

Target:	DPP10
Alternative Name:	DPP10 (DPP10 Products)
Background:	Synonyms: Dipeptidyl peptidase 10, Dipeptidyl peptidase 10 non functional, Dipeptidyl peptidase

IV related protein 3, Dipeptidyl peptidase IV-related protein 3, Dipeptidyl peptidase like protein 2, Dipeptidyl peptidase X, Dipeptidyl peptidase-like protein 2, Dipeptidylpeptidase 10, DPL 2, DPL2, DPP 10, DPP X, Dpp10, DPP10_HUMAN, DPPY, DPRP 3, DPRP-3, DPRP3, Inactive dipeptidyl peptidase 10, KIAA1492.

Background: Dipeptidyl peptidases (DPPs) mediate regulatory activity of their substrates and have been linked to a variety of diseases including type 2 diabetes, obesity and cancer. DPPs have post-proline dipeptidyl aminopeptidase activity, cleaving Xaa-Pro dipeptides from the N-termini of proteins. DPPs can bind specific voltage-gated potassium channels and alter their expression and biophysical properties and may also influence T cells. DPP proteins include DPRP1, DPRP2, DPP3, DPP7, DPP10, DPPX and CD26. DPP10 (dipeptidyl-peptidase 10), also known as DPRP3 (dipeptidyl peptidase IV-related protein 3), DPL2 or DPPY, is a non-functional dipeptidyl peptidase which can bind to the potassium channels KV4.1 and KV4.2. It is a single-pass type II membrane protein expressed in spinal cord, adrenal glands, pancreas and brain tissues and may act as a modulator for cell surface expression and activity of KV4.1 and KV4.2.

Gene ID:

57628

Application Details

Application Notes:	

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

	lI	1:
-	เวทก	111111
	land	11110

Expiry Date:

12 months