antibodies .- online.com







anti-DTX2 antibody (AA 331-430) (Cy5)



Overview

Quantity:	100 μL	
Target:	DTX2	
Binding Specificity:	AA 331-430	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DTX2 antibody is conjugated to Cy5	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DTX2	
Isotype:	IgG	
Predicted Reactivity: Human, Mouse, Rat, Horse, Zebrafish		
Purification:	Purified by Protein A.	

Target Details

Target:	DTX2
Alternative Name:	DTX2 (DTX2 Products)
Background:	Synonyms: Deltex 2, Deltex2, hDTX2, Highly similar to hPMSR6 [H.sapiens], Protein deltex 2,

RING finger protein 58,	RNF58. Zinc ion b	bindina protein.	DTX2 HUMAN.
Till to miger protein oo,	11111 00, 21110 1011 1	on raining proteining	D 17(Z_1101VI) (14.

Background: DTX2 belongs to the Deltex family. It contains one RING-type zinc finger and two WWE domains. DTX2 is a regulator of Notch signaling, a signaling pathway involved in cell-cell communications that regulates a broad spectrum of cell-fate determinations. It probably acts both as a positive and negative regulator of Notch, depending on the developmental and cell context, mediates the antineural activity of Notch, possibly by inhibiting the transcriptional activation mediated by MATCH1. DTX2 also functions as an ubiquitin ligase protein in vitro, suggesting that it may regulate the Notch pathway via some ubiquitin ligase activity. The WWE domains are thought to mediate some protein-protein interaction, and are frequently found in ubiquitin ligases. There are two named isoforms.

Pathways: Notch Signaling

Application Details

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months