



[Go to Product page](#)

Datasheet for ABIN1703896  
**anti-ECAT1 antibody (AA 1-100) (Cy5)**

### Overview

Quantity:	100 µL
Target:	ECAT1 (KHDC3L)
Binding Specificity:	AA 1-100
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ECAT1 antibody is conjugated to Cy5
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Ecat1
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by Protein A.

### Target Details

Target:	ECAT1 (KHDC3L)
Alternative Name:	Ecat1 ( <a href="#">KHDC3L Products</a> )
Background:	Synonyms: C6orf221, Chromosome 6 open reading frame 221, ES cell-associated transcript 1

## Target Details

---

protein, HYDM2, KHD3L\_HUMAN, KHDC3-like protein, KHDC3L.

Background: ECAT1 (ES cell-associated transcript 1 protein) is a 217 amino acid protein that belongs to the KHDC1 family. The ECAT1 protein contains an atypical KH domain with amino acid changes at critical sites, suggesting that it may not bind RNA. Expression of ECAT1 appears to be maximal in germinal vesicle oocytes, it tails off through metaphase II oocytes and is undetectable following the completion of the oocyte to embryo transition. Specifically expressed in the oocytes, recent studies suggest that ECAT1 may function as a regulator of genomic imprinting in the oocyte. Defects in ECAT1 are the cause of hydatidiform mole recurrent type 2 (HYDM2), a disorder characterized by excessive trophoblast development that produces a growing mass of tissue inside the uterus at the beginning of a pregnancy. HYDM2 leads to abnormal pregnancies with no embryo, and cystic degeneration of the chorionic villi.

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months