# antibodies -online.com





## Datasheet for ABIN1704218

## anti-PISD antibody (Cy5)



Go to Product page

$\sim$			
	N/P	r\/I	i⊢₩

Background:

Quantity:	100 μL	
Target:	PISD	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PISD antibody is conjugated to Cy5	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human PISD	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	
Target Details		
Target:	PISD	
Alternative Name:	PISD (PISD Products)	

PISD, PISD\_HUMAN, PSD, PSDC, PSSC.

Synonyms: DJ858B16, dJ858B16.2, DKFZp566G2246, Phosphatidylserine decarboxylase,

Background: Enzymes known as phosphatidylserine decarboxylases (PSDs) catalyze the

Phosphatidylserine decarboxylase beta chain, Phosphatidylserine decarboxylase proenzyme,

### **Target Details**

formation of phosphatidylethanolamine from phosphatidylserine via phosphatidylserine decarboxylation. Type I PSDs contain LGST motifs and are found in bacteria and eukaryotic mitochondria, whereas type II PSDs contain GGST motifs and are found in eukaryotic endomembrane systems. PISD (phosphatidylserine decarboxylase), also known as phosphatidylserine decarboxylase proenzyme, PSDC, PSD, PSSC, DJ858B16, dJ858B16.2 or DKFZp566G2246, is a 408 amino acid a type I phosphatidylserine decarboxylase that localizes to the inner mitochondrial membrane. PISD contains a conserved LGST motif which is cleaved to produce two isoforms known as PISD and PISD \_. PISD is capable of forming a heterodimer and is highly expressed in liver and testis. The gene encoding PISD maps to human chromosome 22q12.2.

Gene ID:

23761

### **Application Details**

Precaution of Use:

Storage:

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Concentration:  $1 \mu g/\mu L$ Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

> This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

12 months **Expiry Date:** 

-20 °C