antibodies .- online.com





anti-APOO antibody (AA 101-198) (Cy5)



()	ve	K\ /		A .
	\cup	1 V/	Щ.	V۷

Quantity:	100 μL	
Target:	APOO	
Binding Specificity:	AA 101-198	
Reactivity:	Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This APOO antibody is conjugated to Cy5	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	en: KLH conjugated synthetic peptide derived from human ApoO/Apolipoprotein O	
Isotype:	IgG	
Cross-Reactivity:	Mouse, Rat	
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	APOO	
Alternative Name:	Apolipoprotein O (APOO Products)	

Target Details	
Background:	Synonyms: ApoO, 0610008C08Rik, 1110019O03Rik, Apolipoprotein O, APOO, APOO_HUMAN, FAM121B, Family with sequence similarity 121B, MGC130105, MGC130106, MGC4825, My025, MYO25, OTTHUMP00000023073, Protein FAM121B, RP23-272D10.2. Background: Apolipoproteins are a family of fatty-acid binding proteins that transport fat through the bloodstream in the form of lipoproteins. ApoO (Apolipoprotein O), also known as FAM121B or My025, is a 198 amino acid single-pass membrane protein that belongs to the apolipoprotein family. Expressed ubiquitously with particularly high expression in diabetic heart tissue, apoO functions to promote the transport of cholesterol from macrophage cells and may be involved in regulatory mechanisms that protect lipid accumulation within the heart. ApoO is present in high density lipoproteins (HDLs) and low density lipoproteins (LDLs) and is secreted by an MTP (microsomal triglyceride transfer protein)-dependent mechanism. Two isoforms of apoO exist due to alternative splicing events.
Gene ID:	79135
Application Details	
Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200

Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months

IF(ICC) 1:50-200

For Research Use only

Restrictions: