



[Go to Product page](#)

Datasheet for ABIN1704230
anti-AGO4 antibody (AA 401-500) (Cy5)

Overview

Quantity:	100 µL
Target:	AGO4 (EIF2C4)
Binding Specificity:	AA 401-500
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AGO4 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Argonaute 4/eIF2C4
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	AGO4 (EIF2C4)
Alternative Name:	Argonaute 4/e2C4 (EIF2C4 Products)
Background:	Synonyms: 5730550L01Rik, Ago 4, Ago4, AGO4_HUMAN, Argonaute4, E 2C 4, e-2C 4, e2C 4,

Target Details

E2c4, eukaryotic translation initiation factor 2C 4, eukaryotic translation initiation factor 2C, hAgo4, KIAA1567, Piwi/Argonaute family protein me2C4, Protein argonaute-4.

Background: Eukaryotic translation initiation factor 2C (eIF2C) proteins (argonaute family) influence RNA interference (RNAi) as components of the RNA-inducible silencing complex (RISC) or microRNA (miRNA)-containing ribonucleoprotein particle (miRNP). Small RNAs, including small interfering RNAs (siRNAs) and miRNAs, can silence target genes through mechanisms that utilize RISC or miRNP particles. eIF2C1 (argonaute 1, AGO1, eIF2C, GERP95, Q99) and Dicer1 play a coordinated role in siRNA-mediated gene silencing. eIF2C2 (Slicer, argonaute 2, AGO2, Q10) is a RISC component that can concentrate in cytoplasmic processing bodies (P-bodies) and catalyze mRNA cleavage. Mammalian P-bodies contain mRNAs and have an association with miRNA-induced translational silencing and siRNA-induced mRNA degradation. Additional eIF2C proteins include eIF2C3 (argonaute 3, AGO3), eIF2C4 (argonaute 4, AGO4) and melF2c5 (mouse argonaute 5).

Gene ID: 192670

Pathways: [Fc-epsilon Receptor Signaling Pathway](#), [Regulatory RNA Pathways](#), [EGFR Signaling Pathway](#), [Neurotrophin Signaling Pathway](#), [Cellular Glucan Metabolic Process](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Handling

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months