



[Go to Product page](#)

Datasheet for ABIN1704433  
**anti-GALNT10 antibody (AA 151-250) (Cy5)**

### Overview

Quantity:	100 µL
Target:	GALNT10
Binding Specificity:	AA 151-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GALNT10 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GALNT10/GalNAc-T10
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Cow,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

### Target Details

Target:	GALNT10
Alternative Name:	GALNT10/GalNAc-T10 ( <a href="#">GALNT10 Products</a> )

## Target Details

---

**Background:** Synonyms: DKFZp586H0623, FLJ00205, FLJ11715, GalNAc T10, GalNAc transferase 10, GalNAc-T10, GalNAcT10, GALNT10, GLT10\_HUMAN, Polypeptide GalNAc transferase 10, Polypeptide N-acetylgalactosaminyltransferase 10, pp GalNAc T10, pp GaNTase 10, pp-GaNTase 10, ppGalNAcT10, ppGaNTase 10, Protein UDP acetylgalactosaminyltransferase 10, Protein-UDP acetylgalactosaminyltransferase 10, UDP GalNAc:polypeptide N acetylgalactosaminyltransferase 10, UDP N acetyl alpha D galactosamine:polypeptide N acetylgalactosaminyltransferase 10, UDP-GalNAc:polypeptide N-acetylgalactosaminyltransferase 10.

**Background:** The UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase (GalNAc-T) family of enzymes are substrate-specific proteins that catalyze the transfer of GalNAc (N-acetylgalactosaminyl) to serine and threonine residues of various proteins, thereby initiating mucin-type O-linked glycosylation in the Golgi apparatus. GalNAc-T10 (Polypeptide N-acetylgalactosaminyltransferase 10), also known as UDP-GalNAc:polypeptide N-acetylgalactosaminyltransferase 10, is a 603 amino acid single-pass type II membrane protein that prefers Muc5Ac and EA2 peptide substrates. The N-terminal domain is involved in substrate binding and manganese coordination, while the C-terminal domain is involved in UDP-Gal binding and catalytic reaction. GalNAc-T10 is widely expressed, with highest levels found in small intestine. There are four isoforms of GalNAc-T10 that are produced as a result of alternative splicing events.

---

**Gene ID:** 55568

## Application Details

---

**Application Notes:** IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

---

**Restrictions:** For Research Use only

## Handling

---

**Format:** Liquid

---

**Concentration:** 1 µg/µL

---

**Buffer:** Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

---

**Preservative:** ProClin

---

## Handling

---

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months