# antibodies -online.com





# Datasheet for ABIN1704477 anti-DMWD antibody (AA 501-600) (Cy5)

# Go to Froduct pag

( )	1 /	0	rv	/ 1 /	71	Α.
	1//	-	1 \/	16		1/1/
$\sim$	v	$\sim$	1 V	١,	_	v v

Quantity:	100 μL	
Target:	DMWD	
Binding Specificity:	AA 501-600	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DMWD antibody is conjugated to Cy5	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human DMWD/DMRN9		
Isotype:	IgG		
Predicted Reactivity:	Human,Mouse,Rat,Cow		
Purification:	Purified by Protein A.		

## **Target Details**

Target:	DMWD
Alternative Name:	DMWD/DMRN9 (DMWD Products)
Background: Synonyms: dystrophia myotonica containing WD repeat mot,D19S593E, DM 9, DM9, DMR 1	

DMR N9 protein, DMRN 9, DMRN9, DMWD, DMWD\_HUMAN, Dystrophia myotonica containing WD repeat mot, Dystrophia myotonica containing WD repeat mot protein, Dystrophia myotonica WD repeat containing protein, Dystrophia myotonica WD repeat-containing protein, Dystrophia myotonica-containing WD repeat mot protein, Gene59, Protein 59, Protein DMR-N9.

Background: DMWD is a 674 amino acid protein containing five WD repeats. DMWD may play a role in the development of mental symptoms in severe cases of myotonic dystrophy, a chronic multisystemic disease characterized by wasting of the muscles, heart conduction defects, cataracts, endocrine changes and myotonia. The DMWD gene is located upstream of the DMPK gene and is prominently expressed in tissues affected in myotonic dystrophy patients. DMWD may also contribute to regulation in meiosis. DMWD is expressed in kidney and spleen, with strongest expression in brain, liver and testis. The gene encoding DMWD maps to human chromosome 19q13.32 and mouse chromosome 7 A3.

Gene ID:

1762

#### **Application Details**

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

## Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	