



[Go to Product page](#)

Datasheet for ABIN1704920
anti-GFM1 antibody (AA 401-500) (Cy5)

Overview

Quantity:	100 µL
Target:	GFM1
Binding Specificity:	AA 401-500
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GFM1 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GFM1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	GFM1
Alternative Name:	GFM1 (GFM1 Products)

Target Details

Background: Synonyms: COXPD1, EF-Gmt, EFG, EFG1, EFGM, EFGM_HUMAN, EGF1, Elongation factor G 1, Elongation factor G 1 mitochondrial, Elongation factor G, Elongation factor G1, FLJ12662, FLJ13632, FLJ20773, G elongation factor mitochondrial 1, G translation elongation factor mitochondrial, GFM 1, GFM, gfm1, hEFG1, mEF G 1, mEF-G 1, mEFG 1, mitochondrial, Mitochondrial elongation factor G1.

Background: Mitochondrial GTPase that catalyzes the GTP-dependent ribosomal translocation step during translation elongation. During this step, the ribosome changes from the pre-translocational (PRE) to the post-translocational (POST) state as the newly formed A-site-bound peptidyl-tRNA and P-site-bound deacylated tRNA move to the P and E sites, respectively. Catalyzes the coordinated movement of the two tRNA molecules, the mRNA and conformational changes in the ribosome. Does not mediate the disassembly of ribosomes from messenger RNA at the termination of mitochondrial protein biosynthesis.

Gene ID: 85476

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months