# antibodies -online.com





Datasheet for ABIN1705009

anti-ANO3 antibody (AA 801-900) (Cy5)

## Go to Product page

_							
0	V	е	r١	/1	е	V	1

Quantity:	100 μL
Target:	ANO3
Binding Specificity:	AA 801-900
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ANO3 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human TMEM16C/Anoctamin 3
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

# **Target Details**

Target:	ANO3	
Alternative Name:	TMEM16C/Anoctamin 3 (ANO3 Products)	
Background: Synonyms: ANO3, C11orf25, GENX 3947, TMEM16C, Transmembrane protein 16C e		

		T 1	160	A N I O O I II I I A A A A I
membrane spannii	na domaine	Iranemamhrana	nrotain Ibi'	$\Delta N U \simeq H I I M \Delta N$
IIICIIIDIANE SPANINI	ių uvinains		protein roo	, AINOS_I IOIVIAIN.

Background: TMEM16C is a 981 amino acid multi-pass membrane protein that is encoded by a gene which maps to chromosome 11. With approximately 135 million base pairs and 1,400 genes, chromosome 11 makes up around 4 % of human genomic DNA and is considered a gene and disease association dense chromosome. The chromosome 11 encoded Atm gene is important for regulation of cell cycle arrest and apoptosis following double strand DNA breaks. Atm mutation leads to the disorder known as ataxia-telangiectasia. The blood disorders Sickle cell anemia and thalassemia are caused by HBB gene mutations, while Wilms' tumors, WAGR syndrome and Denys-Drash syndrome are associated with mutations of the WT1 gene. Jervell and Lange-Nielsen syndrome, Jacobsen syndrome, Niemann-Pick disease, hereditary angioedema and Smith-Lemli-Opitz syndrome are also associated with defects in chromosome 11.

Gene ID:

63982

#### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months