antibodies -online.com





anti-Ethanolamine Kinase 2 antibody (AA 21-120) (Cy5)



Go to Product page

| () | ۱ ۱ | \cap | r | /1 | \cap | ۱ ۸ | 1 |
|-----|-----|--------|----|----|--------|-----|---|
| 0 | 'V | ㄷ | I١ | νı | \Box | V | ۷ |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | Ethanolamine Kinase 2 (ETNK2) |
| Binding Specificity: | AA 21-120 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Ethanolamine Kinase 2 antibody is conjugated to Cy5 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human ETNK2/Ethanolamine kinase 2 |
|-----------------------|---|
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat,Pig |
| Purification: | Purified by Protein A. |

Target Details

| Target: | Ethanolamine Kinase 2 (ETNK2) |
|-------------------|---|
| Alternative Name: | ETNK2/Ethanolamine kinase 2 (ETNK2 Products) |
| Background: | Synonyms: 4933417N20Rik, Al197444, EC 2.7.1.82, EKI 2, EKI2, EKI2_HUMAN, Ethanolamine |

kinase 2, Ethanolamine kinase like protein, Ethanolamine kinase-like protein, ETNK 2, ETNK2, FLJ10761, HMFT1716, RP11 74C13.2, Tuc1.

Background: Ethanolamine kinase 2, also known as EKI2, ETNK2 or HMFT1716, is a 386 amino acid protein that belongs to the choline/ethanolamine kinase family. Via the cytidine diphosphate (CDP) ethanolamine pathway, Ethanolamine kinase 2 catalyses the initial step of phosphatidylethanolamine (PtdEtn) biosynthesis. Ethanolamine kinase 2 is expressed in kidney, liver, testis, ovary and prostate, and is highly specific for ethanolamine phosphorylation. Upregulated during testis development, Ethanolamine kinase 2 may play an essential role in regulating placental hemostasis. Existing as three alternatively spliced isoforms, the gene encoding Ethanolamine kinase 2 maps to human and mouse chromosome 1. Human chromosome 1 spans 260 million base pairs, contains over 3,000 genes, comprises nearly 8 % of the human genome and houses a large number of disease-associated genes, including those that are involved in familial adenomatous polyposis, Stickler syndrome, Parkinson?s disease, Gaucher disease, schizophrenia and Usher syndrome.

Gene ID:

55224

Application Details

| Application | Notes: |
|-------------|--------|

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |

| | lI | 1: |
|---|------|--------|
| - | เวทก | 111111 |
| | land | 11110 |
| | | |

Expiry Date:

12 months