



[Go to Product page](#)

Datasheet for ABIN1705593
anti-DOK7 antibody (AA 21-120) (Cy5.5)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | DOK7 |
| Binding Specificity: | AA 21-120 |
| Reactivity: | Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This DOK7 antibody is conjugated to Cy5.5 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human DOK7 |
| Isotype: | IgG |
| Cross-Reactivity: | Rat |
| Predicted Reactivity: | Human, Mouse, Dog, Cow, Sheep, Pig, Horse |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | DOK7 |
| Alternative Name: | DOK7 (DOK7 Products) |

Target Details

Background: Synonyms: Docking protein 7, DOK 7, DOK7, DOK7_HUMAN, Downstream of tyrosine kinase 7, Protein Dok-7.

Background: The downstream of kinase family (Dok1-7) are members of a class of docking? proteins that include the tyrosine kinase substrates IRS-1 and Cas, which contain multiple tyrosine residues and putative SH2 binding sites. Based on their similarities, the Dok family of proteins can be divided into three subgroups: Dok-1/2/3, Dok-4/5/6 and Dok-7. Through its interaction with muscle-specific receptor kinase (MuSK), Dok-7 is crucial for neuromuscular synaptogenesis and for MuSK activation. Mice lacking Dok-7 do not form neuromuscular synapses nor acetylcholine receptor clusters. Mutations in the Dok-7 gene can cause congenital myasthenic syndromes (CMA) recessively inherited disorders characterized by muscle weakness.

Pathways: [Skeletal Muscle Fiber Development](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months