

[Go to Product page](#)

Datasheet for ABIN1705721
anti-FNDC3A antibody (AA 501-600) (Cy5.5)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | FNDC3A |
| Binding Specificity: | AA 501-600 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This FNDC3A antibody is conjugated to Cy5.5 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human FNDC3A |
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|---|
| Target: | FNDC3A |
| Alternative Name: | FNDC3A (FNDC3A Products) |
| Background: | Synonyms: Fibronectin type III domain containing protein 3a, FNDC3, HUGO, KIAA0970, |

Target Details

FND3A_HUMAN.

Background: FNDC3A is a 1,134 amino acid protein that belongs to the FNDC3 family of proteins. FNDC3A contains an N-terminal proline-rich region, nine fibronectin type-III domains (none of which contain an RGD sequence) and a hydrophobic C-terminal transmembranous helix. Expressed in a wide variety of tissues, FNDC3A localizes to Golgi vesicles and to the developing acrosome of spermatids. FNDC3A is believed to function in glycosaminoglycan and collagen synthesis. In mice, a mutation in the gene encoding FNDC3A causes male sterility due to defective adhesion between Sertoli cells and spermatids in the seminiferous epithelium. This suggests that FNDC3A plays an important role in spermatogenesis, possibly mediating or maintaining the adhesion between Sertoli cells and spermatids.

Gene ID: 22862

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months