antibodies -online.com





Datasheet for ABIN1705817

anti-PRAM1 antibody (AA 551-650) (Cy5.5)



()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

Quantity:	100 μL	
Target:	PRAM1	
Binding Specificity:	AA 551-650	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PRAM1 antibody is conjugated to Cy5.5	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded	
	Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PRAM1
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	PRAM1
Alternative Name:	PRAM1 (PRAM1 Products)

Target Details

Bacl	//	ro	ıın	٨.
Daci	ĸα	ΙU	uu	u.

Synonyms: PML RAR, PML RARA regulated adaptor molecule 1, PML RARA target gene encoding an Adaptor Molecule 1, PML-RARA-regulated adapter molecule 1, PRAM 1, PRAM, PRAM-1, PRAM_HUMAN, PRAM1.

Background: Complete remission of acute promyelocytic leukemia can be achieved by treating patients with retinoic acid, and PML-RAR-a (promyelocytic leukemia-retinoic acid receptor alpha fusion protein) plays a major role in mediating retinoic acid effects in leukemia cells. The retinoic acid-induced gene, PRAM-1 (PML-RAR-a target gene encoding an adaptor molecule 1) encodes an adaptor protein which is expressed and modulated during normal human myelopoiesis. PRAM-1 expression is hindered by expression of PML-RAR-a. The 718 amino acid PRAM-1 protein contains eight N-terminal proline-rich repeats and several proline residues that are clustered as type I or type II SH3 recognition motifs. PRAM-1 demonstrates expression in hematopoietic tissues and lung.

Pathways:

Regulation of Leukocyte Mediated Immunity

Application Details

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months