antibodies .- online.com





anti-ARPC4 antibody (AA 1-100) (Cy5.5)



Overview

Quantity:	100 μL	
Target:	ARPC4	
Binding Specificity:	AA 1-100	
Reactivity:	Xenopus laevis	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ARPC4 antibody is conjugated to Cy5.5	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ARPC4/p20-ARC	
Isotype:	IgG	
Cross-Reactivity:	Xenopus laevis	
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Chicken,Rabbit,Zebrafish	
Purification:	Purified by Protein A.	

Target Details

Target:	ARPC4
Alternative Name:	ARPC4/p20-ARC (ARPC4 Products)

Target Details

Storage Comment:

rarget betane		
Background:	Synonyms: Actin related protein 2/3 complex subunit 4, Actin related protein 2/3 complex	
	subunit 4 20 kD, Actin related protein 2/3 complex subunit 4 20 kDa, Actin-related protein 2/3	
	complex subunit 4, ARC20, Arp2/3 complex 20 kDa subunit, Arp2/3 protein complex subunit	
	p20, ARPC 4, ARPC4, ARPC4_HUMAN, MGC13544, OTTHUMP00000165252,	
	OTTHUMP00000207520, p20 Arc, p20-ARC, p20ARC.	
	Background: This gene encodes one of seven subunits of the human Arp2/3 protein complex.	
	This complex controls actin polymerization in cells and has been conserved throughout	
	eukaryotic evolution. This gene encodes the p20 subunit, which is necessary for actin	
	nucleation and high-affinity binding to F-actin. Alternative splicing results in multiple transcript	
	variants. Naturally occurring read-through transcription exists between this gene and the	
	downstream tubulin tyrosine ligase-like family, member 3 (TTLL3), which results in the	
	production of a fusion protein. [provided by RefSeq, Nov 2010].	
Gene ID:	10093	
Pathways:	RTK Signaling, Regulation of Actin Filament Polymerization	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

	_	
-	II	1:
-	-222	nnc
-	Hand	1111()

Expiry Date:

12 months