# antibodies -online.com





# anti-GNG7 antibody (AA 2-65) (Cy5.5)



#### Overview

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | GNG7   |
| Binding Specificity: | AA 2-65  |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This GNG7 antibody is conjugated to Cy5.5  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human G gamma7 |
|-----------------------|--|
| Isotype:              | IgG  |
| Predicted Reactivity: | Human,Mouse,Rat,Cow,Chicken                                  |
| Purification:         | Purified by Protein A.                                       |

#### **Target Details**

| Target:           | GNG7  |
|-------------------|---|
| Alternative Name: | G gamma7 (GNG7 Products)  |
| Background:       | Synonyms: FLJ00058, GBG7_HUMAN, Gng7, GNGT7, Guanine nucleotide-binding protein |

### Target Details

| GI/GS/GO gamma-7 subunit, Guanine nucleotide-binding protein GI/GS/GO subunit gamma-7.              |  |  |
|---|--|--|
| Background: Guanine nucleotide-binding proteins (G proteins) are involved as a modulator or         |  |  |
| transducer in various transmembrane signaling systems. The beta and gamma chains are                |  |  |
| required for the GTPase activity, for replacement of GDP by GTP, and for G protein-effector         |  |  |
| interaction. Plays a role in the regulation of adenylyl cyclase signaling in certain regions of the |  |  |
| brain. Plays a role in the formation or stabilization of a G protein heterotrimer (G(olf) subunit   |  |  |
| alpha-beta-gamma-7) that is required for adenylyl cyclase activity in the striatum.                 |  |  |
|   |  |  |

Gene ID:

2788

## **Application Details**

| Application Notes: | IF(IHC-P) 1:50-200    |
|--------------------|-----------------------|
|                    | IF(IHC-F) 1:50-200    |
|                    | IF(ICC) 1:50-200      |
| Restrictions:      | For Research Use only |

## Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |