

Datasheet for ABIN1706305
anti-RASA1 antibody (AA 451-550) (Cy5.5)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	RASA1
Binding Specificity:	AA 451-550
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RASA1 antibody is conjugated to Cy5.5
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Ras GTPase-activating protein 1
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Sheep,Pig,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	RASA1
Alternative Name:	RASA1 (RASA1 Products)

Target Details

Background:	Synonyms: Ras GAP, CM AVM, CMAVM, DKFZp434N071, GAP, GTPase activating protein, GTPase-activating protein, OTTHUMP00000222390, OTTHUMP00000222391, OTTHUMP00000222392, OTTHUMP00000222393, p120GAP, p120RASGAP, PKWS, Ras GTPase-activating protein 1, RAS p21 protein activator GTPase activating protein 1, Ras p21 protein activator, RASA, RASA1, RASA1_HUMAN, RasGAP, Triphosphatase activating protein. Background: The mammalian c-H-, c-K- and N-Ras proto-oncogenes encode ubiquitously expressed proteins (1,2). p21Ras can exist in either a physiologically quiescent GDP-binding state or a GTP-binding signal-emitting state (3). Oncogenic p21Ras proteins are trapped in the excited signal-emitting state because the mechanism normally employed to delimit their excitation period, hydrolysis of their bound GTP to GDP, is impaired as a result of specific mutations (3). Interaction of p21Ras with GTPase activating protein (GAP) can increase hydrolysis of p21Ras-bound GTP by as much as 1000-fold (4,5). The product of the neurofibromatosis type 1 gene (NF1) has also been shown to exhibit p21Ras GAP activity (6,7), and proteins that stimulate the GTPase activity of three other low molecular weight GTPases, including Rho, Rab 3A and Rap 1, have also been described (8,9).
Gene ID:	5921
Pathways:	Regulation of Actin Filament Polymerization , Signaling of Hepatocyte Growth Factor Receptor , VEGFR1 Specific Signals

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months