



[Go to Product page](#)

Datasheet for ABIN1706462  
**anti-Golgin A3 antibody (AA 951-1050) (Cy5.5)**

### Overview

Quantity:	100 µL
Target:	Golgin A3 (GOLGA3)
Binding Specificity:	AA 951-1050
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Golgin A3 antibody is conjugated to Cy5.5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Golgin 160
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Cow, Sheep, Rabbit
Purification:	Purified by Protein A.

### Target Details

Target:	Golgin A3 (GOLGA3)
Alternative Name:	Golgin 160 ( <a href="#">GOLGA3 Products</a> )
Background:	Synonyms: GCP170, GOLGA 3, GOLGA3, Golgi autoantigen, Golgi complex associated protein of

## Target Details

---

170 kDa, Golgi complex-associated protein of 170 kD, Golgi membrane associated protein, Golgi peripheral membrane protein, Golgin 160, Golgin subfamily a 3, HGNC:4426, Male enhanced antigen 2, MEA2, SY2/SY10 protein, GOGA3\_HUMAN.

Background: The Golgi apparatus consists of a series of stacked, flattened membrane stacks called cisternae that are involved in the transport of lipids and proteins in the secretory pathway and are important for Golgi-microtubule interaction. Golgin 160 is a 1,498 amino acid protein that localizes to both the cytoplasm and to the Golgi apparatus and contains a series of coiled-coil domains. Expressed in a variety of tissues, including heart, liver, testis, kidney, lung and salivary gland, golgin 160 functions as a homodimer that interacts with GOLGA7 and is thought to be involved in maintaining Golgi structure and may play a role in nuclear transport and Golgi apparatus localization. Multiple isoforms of golgin 160 exist due to alternative splicing events.

---

Gene ID: 2802

Pathways: [SARS-CoV-2 Protein Interactome](#)

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Concentration: 1 µg/µL

---

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

---

Preservative: ProClin

---

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

---

Storage: -20 °C

---

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

---

Expiry Date: 12 months