



[Go to Product page](#)

Datasheet for ABIN1706617
anti-FRAT1 antibody (AA 161-260) (Cy5.5)

Overview

Quantity:	100 µL
Target:	FRAT1
Binding Specificity:	AA 161-260
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FRAT1 antibody is conjugated to Cy5.5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FRAT1
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow
Purification:	Purified by Protein A.

Target Details

Target:	FRAT1
Alternative Name:	FRAT1 (FRAT1 Products)
Background:	Synonyms: FRAT 1, frequently rearranged in advanced T cell lymphomas, Frequently rearranged

Target Details

in advanced T-cell lymphomas, GSK 3 binding protein FRAT1, proto oncogene FRAT1, FRAT1_HUMAN.

Background: FRAT1 and FRAT2 were originally characterized as proteins frequently rearranged in advanced T cell lymphoma, and they have since been identified as proto-oncogenes involved in tumorigenesis. These proteins share significant homology with the Xenopus glycogen synthase kinase-3 (xGSK-3) binding protein, which is designated GBP and is essential for the formation of the dorsal-ventral axis during embryonic development. Establishment of these embryonic axes is mediated by the Wnt intracellular signaling pathway. Wnt signaling is regulated in part by the activity of GSK-3, which phosphorylates and thereby facilitates the degradation of β catenin. GBP binds to GSK-3 and inhibits this phosphorylation, resulting in the accumulation of β catenin and the subsequent transcription of Wnt target genes. Like GBP, FRAT2 has been shown to bind xGSK-3, suggesting that FRAT1 and FRAT2 may be GSK-3 regulatory proteins.

Gene ID: 10023

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 μ g/ μ L

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months