antibodies -online.com





anti-GCP6 antibody (AA 651-750) (Cy5.5)



Go to Product page

| \sim | | | |
|--------|-----|-----|-----|
| | N/P | r\/ | i⊢₩ |

| Quantity: | 100 μL |
|----------------------|---|
| Target: | GCP6 (TUBGCP6) |
| Binding Specificity: | AA 651-750 |
| Reactivity: | Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This GCP6 antibody is conjugated to Cy5.5 |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human GCP6 |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Rat |
| Predicted Reactivity: | Human,Mouse,Dog,Cow,Pig,Horse |
| Purification: | Purified by Protein A. |

Target Details

| Target: | GCP6 (TUBGCP6) |
|-------------------|-------------------------|
| Alternative Name: | GCP6 (TUBGCP6 Products) |

Target Details

| Background: |
|-------------|
|-------------|

Synonyms: GCP 6, TUBGCP6, tubulin, gamma complex associated protein 6, GCP6_HUMAN. Background: The g-Tubulin complex is composed of g Tubulin and the g-Tubulin complex-associated proteins GCP2, GCP3, GCP4, GCP5 and GCP6, all of which are essential components of microtubule organizing centers. g-Tubulin complex components are localized to both the centrosome, where they are involved in microtubule nucleation, and to the cytoplasm, where they exist as soluble complexes that can be recruited to the centrosome as needed. Although the GCP proteins are related, they have distinct roles which contribute to the proper function of the g-Tubulin complex. GCP6 (g-Tubulin complex component 6), also known as TUBGCP6, localizes to the centrosome and is a ubiquitously expressed 1,819 amino acid member of the g-Tubulin complex. Unlike GCP3 and GCP2, GCP6 is not well conserved among eukaryotes. Three isoforms of GCP6 exist due to alternative splicing events.

Gene ID:

85378

Pathways:

M Phase

Application Details

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |