antibodies .- online.com





anti-Ethanolamine Kinase 2 antibody (AA 21-120) (Cy5.5)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
Target:	Ethanolamine Kinase 2 (ETNK2)	
Binding Specificity:	AA 21-120	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Ethanolamine Kinase 2 antibody is conjugated to Cy5.5	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ETNK2/Ethanolamine kinase 2
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Pig
Purification:	Purified by Protein A.

Target Details

Target:	Ethanolamine Kinase 2 (ETNK2)	
Alternative Name:	ETNK2/Ethanolamine kinase 2 (ETNK2 Products)	
Background:	Synonyms: 4933417N20Rik, Al197444, EC 2.7.1.82, EKI 2, EKI2, EKI2_HUMAN, Ethanolamine	

kinase 2, Ethanolamine kinase like protein, Ethanolamine kinase-like protein, ETNK 2, ETNK2, FLJ10761, HMFT1716, RP11 74C13.2, Tuc1.

Background: Ethanolamine kinase 2, also known as EKI2, ETNK2 or HMFT1716, is a 386 amino acid protein that belongs to the choline/ethanolamine kinase family. Via the cytidine diphosphate (CDP) ethanolamine pathway, Ethanolamine kinase 2 catalyses the initial step of phosphatidylethanolamine (PtdEtn) biosynthesis. Ethanolamine kinase 2 is expressed in kidney, liver, testis, ovary and prostate, and is highly specific for ethanolamine phosphorylation. Upregulated during testis development, Ethanolamine kinase 2 may play an essential role in regulating placental hemostasis. Existing as three alternatively spliced isoforms, the gene encoding Ethanolamine kinase 2 maps to human and mouse chromosome 1. Human chromosome 1 spans 260 million base pairs, contains over 3,000 genes, comprises nearly 8 % of the human genome and houses a large number of disease-associated genes, including those that are involved in familial adenomatous polyposis, Stickler syndrome, Parkinson?s disease, Gaucher disease, schizophrenia and Usher syndrome.

Gene ID:

55224

Application Details

Application	Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

\vdash	land	lına
	iaria	шц

Expiry Date:

12 months