



[Go to Product page](#)

Datasheet for ABIN1707119
anti-DNAJB2 antibody (AA 21-120) (Cy5.5)

Overview

Quantity:	100 µL
Target:	DNAJB2
Binding Specificity:	AA 21-120
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DNAJB2 antibody is conjugated to Cy5.5
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HSJ1/DNAJB2
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	DNAJB2
Alternative Name:	HSJ1 (DNAJB2 Products)
Background:	Synonyms: Descriptions, DnaJ Hsp40 homolog subfamily B member 2, DnaJ homolog

Target Details

subfamily B member 2, DnaJ protein homolog 1, DNAJB2, Heat shock 40 kDa protein 3, Heat shock protein J1, Heat shock protein neuronal DNAJ like 1, HSJ 1, HSJ1, HSPF3, DNJB2_HUMAN.

Background: The DnaJ family is one of the largest of all the chaperone families and has evolved with diverse cellular localization and functions. The presence of the J domain defines a protein as a member of the DnaJ family. DnaJ heat shock induced proteins are from the bacterium *Escherichia coli* and are under the control of the htpR regulatory protein. The DnaJ proteins play a critical role in the HSP 70 chaperone machine by interacting with HSP 70 to stimulate ATP hydrolysis. The proteins contain cysteine rich regions that are composed of zinc fingers that form a peptide binding domain responsible for the chaperone function. DnaJ proteins are important mediators of proteolysis and are involved in the regulation of protein degradation, exocytosis and endocytosis. DnaJB2 (DnaJ homolog subfamily B member 2), also known as HSJ1 or HSPF3, is expressed almost exclusively in the brain, with the highest levels in the frontal cortex and hippocampus. Two isoforms are produced due to alternative splicing.

Gene ID: 3300

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months