.-online.com antibodies

Datasheet for ABIN1707197 anti-G Protein-Coupled Receptor 128 antibody (Cy5.5)



Overview

Quantity:	100 µL
Target:	G Protein-Coupled Receptor 128 (GPR128)
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This G Protein-Coupled Receptor 128 antibody is conjugated to Cy5.5
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Western Blotting (WB)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from mouse G protein coupled receptor 128
Isotype:	lgG
Predicted Reactivity:	Mouse,Rat
Purification:	Purified by Protein A.

Target Details

Target:	G Protein-Coupled Receptor 128 (GPR128)
Alternative Name:	Gpr128 (GPR128 Products)
Background:	Synonyms: FLJ14454, G protein coupled receptor 128, GP128_HUMAN, Gpr128, Probable G- protein coupled receptor 128, GPCR GPR128.
	Background: G protein-coupled receptors (GPCRs), also designated seven transmembrane

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1707197 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

(7TM) receptors and heptahelical receptors, are a protein family which interact with G proteins (heterotrimeric GTPases) to synthesize intracellular second messengers such as diacylglycerol,
(hatarotrimaric CTPasas) to synthesize intracellular second messonders such as disculated
cyclic AMP, inositol phosphates, and calcium ions. Their diverse biological functions range from
vision and olfaction to neuronal and endocrine signaling and are involved in many pathological
conditions. G protein receptor 128 (GPR128), a member of the secretin family of GCPRs with a
GPS domain in its N-terminal domain, may mediate signaling processes to the interior of the
cell via activation of G proteins. GPR128 represents an allopeptide which may be involved in T
cell mediated transplant rejection as it is able to stimulate 2.102 T cells.
239853
F(IHC-P) 1:50-200
For Research Use only
Liquid
1 μg/μL
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
50 % Glycerol.
Sodium azide
This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE, which
should be handled by trained staff only.
20 °C
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
12 months