

Datasheet for ABIN1707551 anti-DOK7 antibody (AA 21-120) (Cy7)



Go to Product page

Overview	
Quantity:	100 μL
Target:	DOK7
Binding Specificity:	AA 21-120
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DOK7 antibody is conjugated to Cy7
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Product Details Immunogen:	KLH conjugated synthetic peptide derived from human DOK7
	KLH conjugated synthetic peptide derived from human DOK7
Immunogen:	
Immunogen: Isotype:	IgG
Immunogen: Isotype: Cross-Reactivity:	lgG Rat
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity:	IgG Rat Human,Mouse,Dog,Cow,Sheep,Pig,Horse
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity: Purification:	IgG Rat Human,Mouse,Dog,Cow,Sheep,Pig,Horse

Target Details

Background:	Synonyms: Docking protein 7, DOK 7, DOK7, DOK7_HUMAN, Downstream of tyrosine kinase 7,
	Protein Dok-7.
	Background: The downstream of kinase family (Dok1-7) are members of a class of docking?
	proteins that include the tyrosine kinase substrates IRS-1 and Cas, which contain multiple
	tyrosine residues and putative SH2 binding sites. Based on their similarities, the Dok family of
	proteins can be divided into three subgroups: Dok-1/2/3, Dok-4/5/6 and Dok-7. Through its
	interaction with muscle-specific receptor kinase (MuSK), Dok-7 is crucial for neuromuscular
	synaptogenesis and for MuSK activation. Mice lacking Dok-7 do not form neuromuscular
	synapses nor acetylcholine receptor clusters. Mutations in the Dok-7 gene can cause
	congenital myasthenic syndromes (CMA) recessively inherited disorders characterized by
	muscle weakness.
Pathways:	Skeletal Muscle Fiber Development
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months