
antibodies .- online.com







anti-Leupaxin antibody (AA 251-350) (Cy7)



Overview

Quantity:	100 μL
Target:	Leupaxin (LPXN)
Binding Specificity:	AA 251-350
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Leupaxin antibody is conjugated to Cy7
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Leupaxin
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	Leupaxin (LPXN)
Alternative Name:	Leupaxin (LPXN Products)
Background:	Synonyms: LDLP, LDPL, LPXN, LPXN_HUMAN.

Background: Leupaxin is a 386 amino acid cytoplasmic protein and member of the paxillin family. Leupaxin is highly expressed in lymphoid tissues such as spleen, lymph node, thymus and appendix, with low expression in bone marrow and fetal liver. Consisting of four leucine-rich LD-motifs at the N-terminus and four LIM domains at the C-terminus, leupaxin associates with a member of the focal adhesion kinase family, PYK2, in lymphoid cells. The leupaxin and PYK2 complex is involved in cell type-specific signaling in which it regulates signaling at sites of adhesion. Leupaxin is a substrate for tyrosine kinase in lymphoid cells and is suggested to participate in and be regulated by tyrosine kinase activity. Leupaxin may be a potential progression marker for a subset of prostate cancer and may act as a novel coactivator of the androgen receptor.

Application Details

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200

For Research Use only

Handling

Restrictions:

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months