

Datasheet for ABIN1707877 anti-ARPC4 antibody (AA 1-100) (Cy7)



Go to Product page

Overview		
Quantity:	100 μL	
Target:	ARPC4	
Binding Specificity:	AA 1-100	
Reactivity:	Xenopus laevis	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ARPC4 antibody is conjugated to Cy7	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human ARPC4/p20-ARC	
Isotype:	IgG	
Cross-Reactivity:	Xenopus laevis	
Predicted Reactivity:	Human, Mouse, Rat, Cow, Sheep, Chicken, Rabbit, Zebrafish	
Purification:	Purified by Protein A.	
Target Details		
Target:	ARPC4	

ARPC4/p20-ARC (ARPC4 Products)

Alternative Name:

Target Details

Background:	Synonyms: Actin related protein 2/3 complex subunit 4, Actin related protein 2/3 complex		
	subunit 4 20 kD, Actin related protein 2/3 complex subunit 4 20 kDa, Actin-related protein 2/3		
	complex subunit 4, ARC20, Arp2/3 complex 20 kDa subunit, Arp2/3 protein complex subunit		
	p20, ARPC 4, ARPC4, ARPC4_HUMAN, MGC13544, OTTHUMP00000165252,		
	OTTHUMP00000207520, p20 Arc, p20-ARC, p20ARC.		
	Background: This gene encodes one of seven subunits of the human Arp2/3 protein complex.		
	This complex controls actin polymerization in cells and has been conserved throughout		
	eukaryotic evolution. This gene encodes the p20 subunit, which is necessary for actin		
	nucleation and high-affinity binding to F-actin. Alternative splicing results in multiple transcript		
	variants. Naturally occurring read-through transcription exists between this gene and the		
	downstream tubulin tyrosine ligase-like family, member 3 (TTLL3), which results in the		
	production of a fusion protein. [provided by RefSeq, Nov 2010].		
Gene ID:	10093		
Pathways:	RTK Signaling, Regulation of Actin Filament Polymerization		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		

	1	l:
-	ココロロ	urnen
	Hand	III IU

Expiry Date:

12 months