# antibodies -online.com





## anti-BMP8B antibody (AA 311-402) (Cy7)



Go to	Prod	uct	page

_							
0	V	е	r١	/1	е	V	1

Quantity:	100 μL
Target:	BMP8B (BMP8)
Binding Specificity:	AA 311-402
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BMP8B antibody is conjugated to Cy7
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human BMP8b
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Chicken
Purification:	Purified by Protein A.

#### **Target Details**

Target:	BMP8B (BMP8)
Alternative Name:	BMP8b (BMP8 Products)
Background:	Synonyms: BMP-8, BMP-8B, BMP8, Bmp8b, BMP8B_HUMAN, Bone morphogenetic protein 8,

Bone morphogenetic protein 8B, OP-2, OP2, Osteogenic protein 2.
Background: Bone morphogenic proteins (BMPs) are members of the TGF Beta superfamily.
BMPs are involved in the induction of cartilage and bone formation. In vivo studies have shown
that BMP-2 (also designated BMP-2A) and BMP-3 can independently induce cartilage
formation. Smad3 association with the TGF Beta receptor complex and Smad1 translocation to
the nucleus are observed after the addition of BMP-4 (also designated BMP-2B), suggesting
that BMP-4 may play a role in activation of the Smad pathway. BMP-5, BMP-6 and BMP-7 all
share high sequence homology with BMP-2, indicating that they each may be able to induce
cartilage formation. BMP-8 (also designated OP-2) is thought to be involved in early
development, as detectable expression has not been found in adult organs.

Gene ID:

656

### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200

Restrictions:

For Research Use only

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months