

Datasheet for ABIN1708205 anti-GCHFR antibody (AA 21-60) (Cy7)



Quantity:	100 μL
Target:	GCHFR
Binding Specificity:	AA 21-60
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GCHFR antibody is conjugated to Cy7
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human GCHFR
Isotype:	IgG
Cross-Reactivity:	Rat

Target Details

Purification:

Predicted Reactivity:

Overview

GCHFR Target: Alternative Name: **GCHFR (GCHFR Products)**

Human, Mouse, Cow, Pig, Chicken

Purified by Protein A.

Target Details

HFR, GFRP, GFRP_HUMAN, GTP cyclohydrolase 1 feedback regulatory protein, olase I feedback regulator, GTP cyclohydrolase I feedback regulatory protein, iC138467, MGC138469, p35. TP cyclohydrolase I feedback regulatory protein (GFRP) is encoded by the gene mediates feedback inhibition of GTP cyclohydrolase I activity by terin. GFRP also acts as a mediator for the stimulatory effect of phenylalanine on y. L-phenylalanine reverses this inhibition. The molecular weight of GFRP is
CC138467, MGC138469, p35. TP cyclohydrolase I feedback regulatory protein (GFRP) is encoded by the gene mediates feedback inhibition of GTP cyclohydrolase I activity by terin. GFRP also acts as a mediator for the stimulatory effect of phenylalanine on y. L-phenylalanine reverses this inhibition. The molecular weight of GFRP is
TP cyclohydrolase I feedback regulatory protein (GFRP) is encoded by the gene mediates feedback inhibition of GTP cyclohydrolase I activity by terin. GFRP also acts as a mediator for the stimulatory effect of phenylalanine or y. L-phenylalanine reverses this inhibition. The molecular weight of GFRP is
mediates feedback inhibition of GTP cyclohydrolase I activity by terin. GFRP also acts as a mediator for the stimulatory effect of phenylalanine on y. L-phenylalanine reverses this inhibition. The molecular weight of GFRP is
terin. GFRP also acts as a mediator for the stimulatory effect of phenylalanine on y. L-phenylalanine reverses this inhibition. The molecular weight of GFRP is
y. L-phenylalanine reverses this inhibition. The molecular weight of GFRP is
0. EkDa although aroog linking avnariments have about that CEDD :=
approximately 9.5 kDa although cross-linking experiments have shown that GFRP is usually
nodimer or as a pentamer. 15q15.
200
200
0
Jse only
red solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
ontains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
ned staff only.
Aliquot into multiple vials to avoid repeated freeze-thaw cycles.