antibodies -online.com





anti-GMEB1 antibody (AA 301-400) (Cy7)



Go to Product page

| () | 1/0 | r\ /1 | 014 | |
|-----|-----|-------|-----|---|
| () | ve | I V I | -v | V |
| | | | | |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | GMEB1 |
| Binding Specificity: | AA 301-400 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This GMEB1 antibody is conjugated to Cy7 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human GMEB1 |
|-----------------------|---|
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat,Cow,Sheep,Horse,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | GMEB1 |
|-------------------|---|
| Alternative Name: | GMEB1 (GMEB1 Products) |
| Background: | Synonyms: DNA binding protein p96P, DNA-binding protein p96P, Glucocorticoid modulatory |

element binding protein 1, Glucocorticoid modulatory element-binding protein 1, GMEB 1, GMEB-1, Gmeb1, GMEB1_HUMAN, P96P, Parvovirus initiation factor p96, P 96, P 96, P 96. Background: GMEB-1 is a 573 amino acid protein that contains one SAND domain and is a member of the KDWK family of combinatorial transcription modulators. Localized to both the cytoplasm and the nucleus, GMEB-1 forms a heterodimer with GMEB-2 (Glucocorticoid modulatory element-binding protein 2) and, once associated with GMEB-2, plays a key role in parvovirus DNA replication. In addition, GMEB-1 functions alone as a trans-acting factor that, by binding to glucocorticoid modulatory elements (GMEs) in TAT (tyrosine aminotransferase) promoters, increases intracellular sensitivity to glucocorticoid concentrations. GMEB-1 also interacts with initiator procaspases and, via this interaction, can inhibit caspase-induced apoptosis. Due to alternative splicing events, GMEB-1 is expressed as two isoforms.

Gene ID:

10691

Application Details

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200 IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |