antibodies -online.com





anti-GLUD2 antibody (AA 21-120) (Cy7)



| _ | | |
|-------|------|--|
| ()Ver | view | |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | GLUD2 |
| Binding Specificity: | AA 21-120 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This GLUD2 antibody is conjugated to Cy7 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human GLUD2 |
|-----------------------|---|
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Sheep,Horse,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | GLUD2 |
|-------------------|--|
| Alternative Name: | GLUD2 (GLUD2 Products) |
| Background: | Synonyms: GDH 2, GDH, GDH2, GLUDP1, Glutamate dehydrogenase 2, Glutamate |

dehydrogenase 2 mitochondrial, Glutamate dehydrogenase pseudogene 1.

Background: GLUD2 is both mitochondrial matrix enzymes belonging to the Glu/Leu/Phe/Val dehydrogenases family. Exisiting as homohexamers, GLUD1 catalyzes the oxidative deamination of glutamate to ?ketoglutarate and ammonia while GLUD2 is involved in the recycling of glutamate during neurotransmission. GLUD1 is critical for regulating amino acid induced insulin secretion and is allosterically activated by ADP and inhibited by GTP and ATP. Mutations in the gene encoding GLUD1 causes hyperinsulinism-hyperammonemia syndrome (HHS), which is an inherited condition characterized by high insulin and ammonia levels in the blood. GLUD1 may also be involved in learning and memory reactions by increasing the turnover of the excitatory neurotransmitter glutamate. GLUD2 is expressed in testis and retina, with lower levels found in brain.

Gene ID:

2747

Pathways:

Synaptic Membrane

Application Details

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |