antibodies -online.com





anti-H1F00 antibody (AA 81-180) (Cy7)



Overview

| Quantity: | 100 μL |
|----------------------|---|
| Target: | H1F00 |
| Binding Specificity: | AA 81-180 |
| Reactivity: | Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This H1F00 antibody is conjugated to Cy7 |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human H1F00 |
|-----------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Mouse |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Sheep,Horse,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | H1F00 |
|-------------------|------------------------|
| Alternative Name: | H1F00 (H1F00 Products) |

Target Details

| Background: | Synonyms: H1F00_HUMAN, Histone H1oo, Oocyte-specic histone H1, Oocyte-specic linker |
|---------------------|--|
| | histone H1, osH1, H1F00, H100, OSH1, H1F00 protein. |
| | Background: Histones are basic nuclear proteins that are responsible for the nucleosome |
| | structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 |
| | bp of DNA wrapped around a histone octamer composed of pairs of each of the four core |
| | histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the |
| | interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order |
| | chromatin structures. The protein encoded is a member of the histone H1 family. This gene |
| | contains introns, unlike most histone genes. The protein encoded is a member of the histone |
| | H1 family. The related mouse gene is expressed only in oocytes. [provided by RefSeq, Jul 2008] |
| Gene ID: | 132243 |
| Application Details | |
| Application Notes: | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |
| | |