



[Go to Product page](#)

Datasheet for ABIN1708550
anti-EIF2C3 antibody (AA 251-350) (Cy7)

Overview

Quantity:	100 µL
Target:	EIF2C3
Binding Specificity:	AA 251-350
Reactivity:	Cow, Zebrafish (Danio rerio)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EIF2C3 antibody is conjugated to Cy7
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Argonaute 3/eIF2C3
Isotype:	IgG
Cross-Reactivity:	Cow, Zebrafish (Danio rerio)
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	EIF2C3
Alternative Name:	Argonaute 3/e2C3 (EIF2C3 Products)

Target Details

Background: Synonyms: Argonaute 3/e2C 3, 5730550L01Rik, Ago 3, Ago3, AGO3_HUMAN, argonaute 3, Argonaute3, e 2C 3, e-2C 3, e2C 3, E2c3, e2C3, E2C3 protein, Eukaryotic translation initiation factor 2C 3, Eukaryotic translation initiation factor 2C3, FLJ12765, hAgo3, MGC86946, Protein argonaute-3.

Background: Eukaryotic translation initiation factor 2C (eIF2C) proteins (argonaute family) influence RNA interference (RNAi) as components of the RNA-inducible silencing complex (RISC) or microRNA (miRNA)-containing ribonucleoprotein particle (miRNP). Small RNAs, including small interfering RNAs (siRNAs) and miRNAs, can silence target genes through mechanisms that utilize RISC or miRNP particles. eIF2C1 (argonaute 1, AGO1, eIF2C, GERP95, Q99) and Dicer1 play a coordinated role in siRNA-mediated gene silencing. eIF2C2 (Slicer, argonaute 2, AGO2, Q10) is a RISC component that can concentrate in cytoplasmic processing bodies (P-bodies) and catalyze mRNA cleavage. Mammalian P-bodies contain mRNAs and have an association with miRNA-induced translational silencing and siRNA-induced mRNA degradation. Additional eIF2C proteins include eIF2C3 (argonaute 3, AGO3), eIF2C4 (argonaute 4, AGO4) and melF2c5 (mouse argonaute 5).

Gene ID: 192669

Pathways: [Fc-epsilon Receptor Signaling Pathway](#), [Regulatory RNA Pathways](#), [EGFR Signaling Pathway](#), [Neurotrophin Signaling Pathway](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months