# antibodies -online.com





# anti-BSX antibody (AA 101-200) (Cy7)



Go to Product page

| $\sim$ |    |    |    |             |   |
|--------|----|----|----|-------------|---|
| ()     | VE | ۲۱ | /1 | $\triangle$ | Λ |

| Quantity:            | 100 μL  |
|----------------------|---|
| Target:              | BSX   |
| Binding Specificity: | AA 101-200  |
| Reactivity:          | Mouse, Rat  |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This BSX antibody is conjugated to Cy7  |
| Application:         | Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human BSX |  |
|-----------------------|---|--|
| Isotype:              | IgG   |  |
| Cross-Reactivity:     | Mouse, Rat  |  |
| Predicted Reactivity: | Human,Dog,Cow,Sheep,Pig,Horse,Rabbit                    |  |
| Purification:         | Purified by Protein A.                                  |  |
|                       |   |  |

## **Target Details**

| Target:           | BSX                |
|-------------------|--------------------|
| Alternative Name: | BSX (BSX Products) |

## **Target Details**

| _                   |   |  |  |
|---------------------|---|--|--|
| Background:         | Synonyms: Brain specic homeobox, Brain specic homeobox protein homolog, BSH, BSX1,                  |  |  |
|                     | BSH_HUMAN.  |  |  |
|                     | Background: BSX(Brain-specific homeobox protein homolog) may play a role in the                     |  |  |
|                     | determination and function of cell types in the brain. In brain, it is restricted to a few specific |  |  |
|                     | developing brain structures such as pineal gland, telencephalic septum, hypothalamic pre-           |  |  |
|                     | mammillary body and arcuate nucleus. Bsx might be considered an important molecular                 |  |  |
|                     | marker for early embryonic stages of epiphysis development. It belongs to the distal-less           |  |  |
|                     | homeobox family and contains 1 homeobox DNA-binding domain.   |  |  |
| Gene ID:            | 390259  |  |  |
| Pathways:           | Feeding Behaviour   |  |  |
| Application Details |   |  |  |
| Application Notes:  | FCM 1:20-100  |  |  |
|                     | IF(IHC-P) 1:50-200  |  |  |
|                     | IF(IHC-F) 1:50-200  |  |  |
|                     | IF(ICC) 1:50-200  |  |  |
| Restrictions:       | For Research Use only   |  |  |
| Handling            |   |  |  |
| Format:             | Liquid  |  |  |
| Concentration:      | 1 μg/μL   |  |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and         |  |  |
|                     | 50 % Glycerol.  |  |  |
| Preservative:       | ProClin   |  |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be                 |  |  |
|                     | handled by trained staff only.  |  |  |
| Storage:            | -20 °C  |  |  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                   |  |  |
| Expiry Date:        | 12 months   |  |  |
|                     |   |  |  |