antibodies -online.com







Overview

Quantity:	100 μL
Target:	FBXO9
Binding Specificity:	AA 301-400
Reactivity:	Xenopus laevis
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FBXO9 antibody is conjugated to Cy7
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FBX09
Isotype:	lgG
Cross-Reactivity:	Xenopus laevis
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Rabbit, Zebrafish, Chimpanzee
Purification:	Purified by Protein A.

Target Details

Target:	FBXO9
Alternative Name:	FBX09 (FBX09 Products)

Target Details

Background:

Synonyms: Cross-immune reaction antigen 1, dJ341E18.2, DKFZp434C0118, F-box only protein 9, Fbp24, FBX9, FBX9_HUMAN, Fbxo9, KIAA0936, NY REN 57, OTTHUMP00000016628, OTTHUMP00000016629, OTTHUMP00000214588, Renal carcinoma antigen NY-REN-57, VCIA1.

Background: This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of the ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbxs class. Alternative splicing of this gene generates at least 3 transcript variants diverging at the 5' terminus. [provided by RefSeq, Jul 2008].

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months