# antibodies .- online.com





# anti-GNL2 antibody (AA 1-100) (Cy7)



$\sim$	4.4	D		page
1-()	10	Pron	ויאווו	nane

$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL
Target:	GNL2
Binding Specificity:	AA 1-100
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GNL2 antibody is conjugated to Cy7
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human GNL2
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

## **Target Details**

Target:	GNL2
Alternative Name:	GNL2 (GNL2 Products)

## **Target Details**

Background:	Synonyms: Autoantigen NGP-1, Autoantigen NGP1, DJ423B22.6 novel nucleolar guanosine 5'-
	triphosphate binding protein, FLJ40906, GNL2, Guanine nucleotide binding protein-like 2
	nucleolar, HUMAUANTIG, NGP1, NOG2_HUMAN, Nucleolar GTP binding protein 2, Nucleolar
	GTP-binding protein 2, Nucleolar GTPase.
	Background: GNL2 is a nucleolar guanasine-triphosphate binding protein that is ubiquitously
	expressed at low levels in almost all tissues. GNL2 is involved in the crucial process of
	trafficking proteins out of the nucleus. Specifically, it is a GTPase that interacts with the 60s
	preribosomal subunit in the nucleus and facilitates export of the subunit into the cytoplasm.
	GTPases are responsible for the hydrolysis of GTP by way of a protein region dubbed the G
	domain. GTPases are often involved in the translocating proteins through membranes gleaning
	energy for the activity by hydrolizing GTP. GNL2 shares G domain homology and some
	functionality with nucleostemin (GNL3), another nuclear GTPase. Highest expression of GNL2
	is found in testis.
Gene ID:	29889
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months