# antibodies -online.com





## anti-GNPDA2 antibody (AA 101-200) (Cy7)



#### Overview

Quantity:	100 μL
Target:	GNPDA2
Binding Specificity:	AA 101-200
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GNPDA2 antibody is conjugated to Cy7
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human GNPDA2
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

### Target Details

Target:	GNPDA2
Alternative Name:	GNPDA2 (GNPDA2 Products)

#### **Target Details**

Storage Comment:

Expiry Date:

Target Details	
Background:	Synonyms: GlcN6P deaminase 2, Glucosamine 6 phosphate deaminase 2, Glucosamine 6
	phosphate isomerase 2, Glucosamine 6 phosphate isomerase SB52, Glucosamine-6-phosphate
	deaminase 2, Glucosamine-6-phosphate isomerase 2, Glucosamine-6-phosphate isomerase
	SB52, GNP2, GNPDA 2, Gnpda2, GNPI2_HUMAN, SB52.
	Background: During fertilization in mammals, the sperm activates the egg by causing an
	increase in the level of free cytoplasmic calcium concentration. This increased calcium
	concentration induces a characteristic series of oscillations that trigger egg activation and early
	embryo development. A hamster protein named oscillin is thought to be involved in this
	pathway. The enzyme glucosamine-6-phosphate isomerase (GNPI) or deaminase (GNPDA1)
	and the related protein GNPDA2 are the human homologs of hamster oscillin. GNPDA1 and
	GNPDA2 catalyze the conversion of GNP to fructose-6-phosphate and ammonia. Both proteins
	exist as homohexamers and are ubiquitously expressed with highest expression in testis, ovary
	and heart. Three isoforms of GNPDA2 are expressed due to alternative splicing events.
Gene ID:	132789
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

12 months

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.