# antibodies - online.com







# anti-GDI1 antibody (AA 1-100) (Cy7)



$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL
Target:	GDI1
Binding Specificity:	AA 1-100
Reactivity:	Mouse, Chimpanzee
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GDI1 antibody is conjugated to Cy7
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human GDI1	
Isotype:	lgG	
Cross-Reactivity:	Chimpanzee, Mouse	
Predicted Reactivity:	Human,Rat,Cow,Sheep,Pig,Rabbit,Zebrafish	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	GDI1
Alternative Name:	GDI1 (GDI1 Products)

### **Target Details**

Storage Comment:

l arget Details		
Background:	Synonyms: 1A, FLJ41411, GDI-1, gdi1, GDIA_HUMAN, GDIL, GDP dissociation inhibitor 1,	
	Guanosine diphosphate dissociation inhibitor 1, Mental retardation X-linked 41, Mental	
	retardation X-linked 48, MRX41, MRX48, Oligophrenin 2, Oligophrenin-2, OPHN2, Protein XAP 4,	
	Protein XAP-4, Protein XAP4, Rab GDI alpha, Rab GDP dissociation inhibitor alpha, RABGD1A,	
	RABGDIA, XAP 4.	
	Background: Rab proteins, a family of Ras-related small GTP-binding proteins, play a key role in	
	regulating intracellular vesicle trafficking (1?). Rab GDP dissociation inhibitor (Rab GDI or GDI2)	
	forms a soluble complex with Rab proteins and thereby prevents the exchange of GDP for GTP	
	(1?). In mammals, there exist two major isoforms, Rab GDI alpha (also known as XAP-4) and	
	Rab GDI beta (1,4). While the mammalian Rab GDI beta-genes are ubiquitously expressed, the	
	Rab GDI ?genes are predominantly brain-specific (1). Since it is expressed predominantly in	
	neural and sensory tissues, Rab GDI a may serve a specific function in neural signal	
	transmission (5). The gene sequences for the Rab GDI proteins are extremely conserved in	
	evolution, with substantial homology preserved across three eukaryotic kingdoms (5).	
Gene ID:	2664	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

	lI	1:
-	เวทก	111111
	land	HHU

Expiry Date:

12 months