# antibodies -online.com





# anti-LTK antibody (AA 41-140) (FITC)



Go to Product page

( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL
Target:	LTK
Binding Specificity:	AA 41-140
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LTK antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human LTK
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Horse,Rabbit
Purification:	Purified by Protein A.

#### **Target Details**

Target:	LTK
Alternative Name:	Ltk (LTK Products)
Background:	Synonyms: ALK, Anaplastic lymphoma kinase, Leukocyte receptor tyrosine kinase, Leukocyte

tyrosine kinase, Leukocyte tyrosine kinase rece	eptor, Leukocyte tyrosine kinase receptor, Ltk,
LTK HUMAN, Protein tyrosine kinase 1, TYK 1,	TYK1.

Background: LTK is a member of the ros/insulin receptor family of tyrosine kinases. Tyrosine-specific phosphorylation of proteins is a key to the control of diverse pathways leading to cell growth and differentiation. Multiple transcript variants encoding different isoforms have been found for this gene.

Gene ID: 4058

Pathways: RTK Signaling

### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months