



[Go to Product page](#)

Datasheet for ABIN1709548
anti-IFI30 antibody (AA 181-250) (FITC)

Overview

Quantity:	100 µL
Target:	IFI30
Binding Specificity:	AA 181-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IFI30 antibody is conjugated to FITC
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human IFI30
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat
Purification:	Purified by Protein A.

Target Details

Target:	IFI30
Alternative Name:	I30 (IFI30 Products)
Background:	Synonyms: Gamma interferon inducible lysosomal thiol reductase, Gamma interferon inducible

Target Details

protein IP 30, GILT, I 30, interferon gamma inducible protein 30, Interferon gamma inducible protein 30 preproprotein, IP30, Lysosomal thiol reductase, Lysosomal thiol reductase, gamma-interferon-inducible.

Background: Proteins internalized into the endocytic pathway are usually degraded. Efficient proteolysis requires denaturation, induced by acidic conditions within lysosomes, and reduction of inter- and intrachain disulfide bonds. Cytosolic reduction is mediated enzymatically by thioredoxin. In the endocytic pathway, reduction of protein disulfide bonds is important for the generation of MHC class II-peptide complexes. This process is catalyzed by a gamma-interferon-inducible thiol reductase (GILT). GILT is synthesized as a precursor, and following delivery to MHC class II-containing compartments (MIICs), is processed to the mature form via cleavage of amino- and carboxy-terminal propeptides. A lysosomal thiol reductase, GILT, is optimally active at low pH and capable of catalyzing disulfide bond reduction both in vivo and in vitro. GILT is expressed constitutively in antigen-presenting cells and is induced by g-interferon in other cell types, suggesting a potentially important role in antigen processing. Additionally, T cell recognition of select exogenous and endogenous epitopes is dependent on tumor cell expression of GILT. The absence of GILT in melanomas alters antigen processing and the hierarchy of immunodominant epitope presentation.

Gene ID: 10437

UniProt: [P13284](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months