

[Go to Product page](#)

Datasheet for ABIN1709558

anti-Leupaxin antibody (AA 251-350) (FITC)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | Leupaxin (LPXN) |
| Binding Specificity: | AA 251-350 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Leupaxin antibody is conjugated to FITC |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human Leupaxin |
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Pig,Horse |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | Leupaxin (LPXN) |
| Alternative Name: | Leupaxin (LPXN Products) |
| Background: | Synonyms: LDLP, LDPL, LPXN, LPXN_HUMAN. |

Target Details

Background: Leupaxin is a 386 amino acid cytoplasmic protein and member of the paxillin family. Leupaxin is highly expressed in lymphoid tissues such as spleen, lymph node, thymus and appendix, with low expression in bone marrow and fetal liver. Consisting of four leucine-rich LD-motifs at the N-terminus and four LIM domains at the C-terminus, leupaxin associates with a member of the focal adhesion kinase family, PYK2, in lymphoid cells. The leupaxin and PYK2 complex is involved in cell type-specific signaling in which it regulates signaling at sites of adhesion. Leupaxin is a substrate for tyrosine kinase in lymphoid cells and is suggested to participate in and be regulated by tyrosine kinase activity. Leupaxin may be a potential progression marker for a subset of prostate cancer and may act as a novel coactivator of the androgen receptor.

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months