



[Go to Product page](#)

Datasheet for ABIN1709627

anti-UCMA antibody (AA 21-120) (FITC)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | UCMA |
| Binding Specificity: | AA 21-120 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This UCMA antibody is conjugated to FITC |
| Application: | Western Blotting (WB) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human UCMA |
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|---|
| Target: | UCMA |
| Alternative Name: | UCMA (UCMA Products) |
| Background: | Synonyms: C10orf49, Gla rich protein, Gla-rich protein, Grp, GRP, OTTMUSP00000011599, RP23-272I15.1, UCMA, Ucm-C, UCMA_HUMAN, Unique cartilage matrix associated protein, |

Target Details

Unique cartilage matrix-associated protein C-terminal fragment, Upper zone of growth plate and cartilage matrix associated.

Background: UCMA is a 138 amino acid secreted protein that is highly expressed in resting chondrocytes in developing long bones and is thought to function in the early phase of chondrocyte differentiation. A furin-like protease processes UCMA into an N-terminal 37 amino acid peptide and a C-terminal 74 amino acid peptide, which is referred to as Unique cartilage matrix-associated protein C-terminal fragment (Ucma-C). Introduction of recombinant Ucma-C interferes with osteogenic differentiation of mesenchymal stem cells, MC3T3-E1 preosteoblasts and primary osteoblasts. This suggests that Ucma may be involved in the negative regulation of osteogenic differentiation of osteochondrogenic precursor cells at the cartilage-bone interface and in peripheral zones of fetal cartilage.

Gene ID: 221044

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months