antibodies - online.com







anti-C9orf25 antibody (AA 1-100) (FITC)



/ N	11/0	K /	\sim	A /
	ve	1 \/	ı⊢۱	ΛI
\sim	' V C	1 V		v v

Quantity:	100 μL	
Target:	C9orf25	
Binding Specificity:	AA 1-100	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This C9orf25 antibody is conjugated to FITC	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human C9orf25
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Sheep, Horse, Chicken
Purification:	Purified by Protein A.

Target Details

Target:	C9orf25	
Alternative Name:	C9orf25 (C9orf25 Products)	
Background:	und: Synonyms: bA573M23.5, C9orf25, Chromosome 9 open reading frame 25, F219A_HUMAN	

FLJ39031, Hypothetical protein LOC203259, Uncharacterized protein C9orf25.

Background: C9orf25 (chromosome 9 open reading frame 25) is a 185 amino acid protein that exists as seven alternatively spliced isoforms that are encoded by a gene located on human chromosome 9. Chromosome 9 consists of about 145 million bases, represents 4 % of the human genome and encodes nearly 900 genes. Thought to play a role in gender determination, deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 though through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation leading to the aberrant production of BCR-ABL fusion protein often found in leukemias.

Gene ID:

203259

Application Details

Application Notes:	IF(IHC-P) 1:50-200
--------------------	--------------------

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months