antibodies .- online.com





Datasheet for ABIN1709759

anti-GMEB2 antibody (AA 101-200) (FITC)



Overview

Quantity:	100 μL
Target:	GMEB2
Binding Specificity:	AA 101-200
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GMEB2 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GMEB2
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	GMEB2
Alternative Name:	GMEB2 (GMEB2 Products)

Target Details

rarget Details		
Background:	Synonyms: GMEB2, DNA binding protein p79P, DNA-binding protein p79P, Glucocorticoid	
	modulatory element binding protein 2, Glucocorticoid modulatory element-binding protein 2,	
	GMEB 2, GMEB-2, Gmeb2, GMEB2_HUMAN, KIAA1269, P79P, Parvovirus initiation factor p79, P	
	p79, P79, RP4 663D7.1.	
	Background: GMEB-2 is a DNA-binding protein that plays a crucial role modulating transcription	
	upon activation by steroid hormones. GMEB-2 is ubiquitously expressed with preferential expression in developmentally important tissues, such as testis, bone marrow, placenta and	
	fetal tissues. It localizes to the nucleus and cytoplasm and contains a SAND domain near its N-	
	terminus and a C-terminal coiled coil structure. GMEB-2 functions as a homodimer or as a	
	heterodimer with GMEB-1. The formed complex specifically binds to glucocorticoid modulatory	
	elements (GME) in the promoter region of target genes and recruits the histone acetylase CREB	
	binding protein (CBP) during glucocorticoid signaling. This acts to increase sensitivity to low	
	concentrations of glucocorticoids. In addition, GMEB-2 functions as an auxiliary factor required	
	for parvovirus replication.	
Gene ID:	26205	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

-20 °C

Storage:

Storage Comment:

		1 .	
ш	land	linc	٩
	ıanıu	11111	J

Expiry Date:

12 months