## antibodies -online.com





Go to Product page

## Datasheet for ABIN1709787

## anti-CRTAP antibody (AA 201-300) (FITC)

Overview	
Quantity:	100 μL
Target:	CRTAP
Binding Specificity:	AA 201-300
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CRTAP antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human CRTAP
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.
Target Details	
Target:	CRTAP
Alternative Name:	CRTAP (CRTAP Products)

## **Target Details**

•	
Background:	Synonyms: Cartilage associated protein, LEPREL3, leprecan-like 3, CASP, CRTAP_HUMAN.
	Background: CRTAP is a secreted protein localizing to the extracellular space that plays a role in
	collagen post-translational modifications, extracellular fibril assembly and intracellular
	trafficking. CRTAP is widely expressed with predominant expression in articular chondrocytes.
	It contains a signal peptide and a tetratricopeptide-like helical domain and is essential for
	normal bone formation. In the endoplasmic reticulum (ER), CRTAP forms a complex with Gros1
	and CyPB (cyclophilin B) and is required for the efficient 3-hydroxylation of target prolyl residues
	in Collagen Type I molecules, the major structural proteins of skin and bone. Mutations in the
	gene encoding CRTAP can lead to autosomal recessive osteogenesis imperfecta (OI) type 7
	and type 2B. OI, also known as brittle bone disease, is characterized by bone fragility and
	susceptibility to fractures. OI type 7 is a mild form of this disorder, while OI type 2B is a neonatal
	lethal condition.
Gene ID:	10491
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months