# antibodies - online.com







# anti-APOO antibody (AA 101-198) (FITC)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL
Target:	APOO
Binding Specificity:	AA 101-198
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This APOO antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human ApoO/Apolipoprotein O	
Isotype:	IgG	
Cross-Reactivity:	Mouse, Rat	
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	

# **Target Details**

Target:	APOO	
Alternative Name:	Apolipoprotein O (APOO Products)	

### **Target Details**

Background:	Synonyms: ApoO, 0610008C08Rik, 1110019003Rik, Apolipoprotein O, APOO, APOO_HUMAN,
	FAM121B, Family with sequence similarity 121B, MGC130105, MGC130106, MGC4825, My025,
	MYO25, OTTHUMP00000023073, Protein FAM121B, RP23-272D10.2.
	Background: Apolipoproteins are a family of fatty-acid binding proteins that transport fat
	through the bloodstream in the form of linoproteins. Apon (Apolipoprotein 0), also known as

ransport fat through the bloodstream in the form of lipoproteins. ApoO (Apolipoprotein O), also known as FAM121B or My025, is a 198 amino acid single-pass membrane protein that belongs to the apolipoprotein family. Expressed ubiquitously with particularly high expression in diabetic heart tissue, apoO functions to promote the transport of cholesterol from macrophage cells and may be involved in regulatory mechanisms that protect lipid accumulation within the heart. ApoO is present in high density lipoproteins (HDLs) and low density lipoproteins (LDLs) and is secreted by an MTP (microsomal triglyceride transfer protein)-dependent mechanism. Two isoforms of apoO exist due to alternative splicing events.

Gene ID:

79135

#### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months