antibodies -online.com





anti-GLT25D2 antibody (AA 501-600) (FITC)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	GLT25D2
Binding Specificity:	AA 501-600
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GLT25D2 antibody is conjugated to FITC
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GLT25D2
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Cow,Sheep,Pig,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	GLT25D2
Alternative Name:	GLT25D2 (GLT25D2 Products)

Target Details

Expiry Date:

Background:	Synonyms: C1orf17, Glt25d2, Glycosyltransferase 25 family member 2, GT252_HUMAN,	
	Hydroxylysine galactosyltransferase 2, Procollagen galactosyltransferase 2, RP11-498P10.2.	
	Background: GLT25D2 is a 626 amino acid protein that localizes to endoplasmic reticulum	
	lumen and belongs to the glycosyltransferase 25 family. Possessing beta-galactosyltransferase	
	activity, GLT25D2 transports beta-galactose to hydroxylysine residues on collagen but lacks	
	glucosyltransferase activity. GLT25D2 is expressed in skeletal muscle and brain, and is	
	encoded by a gene located on human chromosome 1q25.3, a region associated with the	
	hereditary prostate cancer (HPC1) locus. Human chromosome 1 spans 260 million base pairs,	
	contains over 3,000 genes, comprises nearly 8 % of the human genome and houses a large	
	number of disease-associated genes, including those that are involved in familial adenomatous	
	polyposis, Stickler syndrome, Parkinson?s disease, Gaucher disease, schizophrenia and Usher	
	syndrome.	
Gene ID:	23127	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

12 months