# antibodies -online.com





#### Go to Product page

## Datasheet for ABIN1710027

## anti-ZFAND5 antibody (AA 121-213) (FITC)

Overview	
Quantity:	100 μL
Target:	ZFAND5
Binding Specificity:	AA 121-213
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZFAND5 antibody is conjugated to FITC
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human ZFAND5

Immunogen:	KLH conjugated synthetic peptide derived from human ZFAND5
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Cow, Sheep, Pig, Chicken, Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	ZFAND5
Alternative Name:	ZFAND5 (ZFAND5 Products)
Background:	Synonyms: ZA20D2, ZNF216, ZFAND5A, AN1-type zinc finger protein 5, Zinc finger A20 domain-

containing protein 2, Zinc finger protein 216, ZFAND5	containing	protein 2.	. Zinc finger	protein 216	. ZFAND5
---	------------	------------	---------------	-------------	----------

Background: Involved in protein degradation via the ubiquitin-proteasome system. May act by anchoring ubiquitinated proteins to the proteasome. Plays a role in ubiquitin-mediated protein degradation during muscle atrophy. Plays a role in the regulation of NF-kappa-B activation and apoptosis. Inhibits NF-kappa-B activation triggered by overexpression of RIPK1 and TRAF6 but not of RELA. Inhibits also tumor necrosis factor (TNF), IL-1 and TLR4-induced NF-kappa-B activation in a dose-dependent manner. Overexpression sensitizes cells to TNF-induced apoptosis. Is a potent inhibitory factor for osteoclast differentiation.

Gene ID:	7763
UniProt:	076080

Pathways: Platelet-derived growth Factor Receptor Signaling

### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months