antibodies .-online.com

Datasheet for ABIN1710063 anti-TMC8 antibody (AA 301-400) (FITC)



\sim						
()	۱/	0	r\	/1	\cap	۱۸/
U	v	C	1	<i>v</i> I	C	W

Quantity:	100 µL	
Target:	TMC8	
Binding Specificity:	AA 301-400	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TMC8 antibody is conjugated to FITC	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human EVER2/TMC8
lsotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Horse
Purification:	Purified by Protein A.

Target Details

Target:	TMC8
Alternative Name:	EVER2/TMC8 (TMC8 Products)
Background:	Synonyms: epidermodysplasia verrucormis 2, Epidermodysplasia verrucormis protein 2, EV2,

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1710063 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

	EVIN2, FLJ40668, FLJ43684, MGC102701, MGC40121, TMC8, Transmembrane channel like
	protein 8, TMC8_HUMAN.
	Background: Transmembrane channel-like protein 8 (TMC8), also known as Epidermodysplasia
	verruciformis protein 2 (EVER2), is a 726 amino acid member of the TMC family of proteins.
	Localized to the endoplasmic reticulum membrane, TMC8 is thought to form a transmembrane
	channel-like protein with eight predicted transmembrane domains and three leucine zipper
	motifs. Mutations in the genes encoding TMC8 and TMC6, another member of the TMC family,
	have been shown to cause epidermodysplasia verruciformis (EV), an autosomal recessive
	dermatosis characterized by abnormal susceptibility to human papillomaviruses (HPVs) and a
	high rate of progression to squamous cell carcinoma on sun-exposed skin. Infection by HPVs
	lead to persistent wart-like or macular lesions. TMC8 is expressed in placenta, prostate and
	testis, and three named isoforms exist as a result of alternative splicing events.
Gene ID:	147138
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN1710063 | 03/07/2024 | Copyright antibodies-online. All rights reserved.