# antibodies - online.com







# anti-Cyclin M3 antibody (AA 251-350) (FITC)



| ( ) | 1/0 | r\ /1 | 014 |   |
|-----|-----|-------|-----|---|
| ( ) | ve  | I V I | -v  | V |
|     |     |       |     |   |

| Quantity:            | 100 μL   |  |
|----------------------|--|--|
| Target:              | Cyclin M3 (CNNM3)  |  |
| Binding Specificity: | AA 251-350   |  |
| Reactivity:          | Human  |  |
| Host:                | Rabbit   |  |
| Clonality:           | Polyclonal   |  |
| Conjugate:           | This Cyclin M3 antibody is conjugated to FITC  |  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |

### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human Cyclin M3 |  |
|-----------------------|---|--|
| Isotype:              | IgG   |  |
| Predicted Reactivity: | Human,Mouse,Rat,Dog   |  |
| Purification:         | Purified by Protein A.  |  |

# **Target Details**

| Target:   | Cyclin M3 (CNNM3)          |  |
|---|----------------------------|--|
| Alternative Name:   | Cyclin M3 (CNNM3 Products) |  |
| ackground: Synonyms: ACDP3, Ancient conserved domain containing protein 3, CNNM3, DKFZp434I10 |                            |  |

| FLJ20018, Metal transporter CNNM3, CNNM3_HUMAN.               |
|---|
| Background: Cyclin M3 is a 707 amino acid multi-pass membrane |

Background: Cyclin M3 is a 707 amino acid multi-pass membrane protein that shares weak sequence similarity with cyclin proteins, yet displays no cyclin-like function in vivo. Though ubiquitously expressed, Cyclin M3 is found at highest levels in kidney, brain, spleen and heart. Cyclin M3 is localized to the nucleus where it is likely a metal transporter. Cyclin M3 contains two CBS domains, which appear to bind ligands with an adenosyl group such as AMP, ATP and S-AdoMet and may play a regulatory role in sensitizing proteins to adenosyl-carrying ligands. There are three isoforms of Cyclin M3 that are produced as a result of alternative splicing events.

Gene ID:

26505

## **Application Details**

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|--------------------|
|                    | IF(IHC-F) 1:50-200 |

IF(ICC) 1:50-200

Restrictions: For Research Use only

### Handling

| Format:            | Liquid   |  |
|--------------------|--|--|
| Concentration:     | 1 μg/μL  |  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |  |
| Preservative:      | ProClin  |  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |  |
| Storage:           | -20 °C   |  |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |  |
| Expiry Date:       | 12 months  |  |